

PP-DG-US-3210 07/2020 ©Lilly USA, LLC 2020. All rights reserved.




Frame 1



Super: One week is 604,800 seconds.

Frame 2



Super: That's 10,080 minutes.

Make every moment precious.

Super: Make every moment precious.

Frame 4

Frame 3



Super: Make every moment count.

Frame 5



Super: Trulicity may help lower A1C*

Frame 6



Super: Once-weekly Trulicity works 24/7.

Frame 7



Super: For all those moments.

Talk to a doctor or visit Trulicity.com to learn more.



PURPOSE AND SAFETY SUMMARY WITH WARNINGS

Important Facts About Trulicity® (Trū-li-si-tee). It is also known as dulaqlutide.

Trulicity is a prescription medicine for adults with type 2 diabetes used to improve blood sugar (glucose) and used to reduce the risk of major cardiovascular events (problems having to do with the heart and blood vessels) such as death, heart attack, or stroke in people who have heart disease or multiple cardiovascular risk factors.

 Trulicity is given through an injection (needle). You take it once a week by injecting it under the skin of your stomach, thigh, or upper arm. Use Trulicity together with the diet and exercise that your doctor recommends. Trulicity is not insulin.

Warning

Trulicity may cause tumors in the thyroid, including thyroid cancer. Watch for possible symptoms, such as a lump or swelling in the neck, trouble swallowing, hoarseness, or shortness of breath. If you have a symptom, tell your doctor.

- Do not use Trulicity if you or any of your family have ever had a type of thyroid cancer called medullary thyroid carcinoma (MTC).
- Do not use Trulicity if you have Multiple Endocrine Neoplasia syndrome type 2 (MEN 2).
- Do not use Trulicity if you are allergic to dulaglutide or other ingredients in Trulicity.

Ask your doctor how to recognize the serious side effects below and what to do if you think you have one:

Inflamed pancreas (pancreatitis). Stop using Trulicity and call your healthcare provider right away if you have severe pain in your stomach area (abdomen), with or without vomiting, that will not go away. You may feel the pain from your abdomen to your back.

Changes in vision. Tell your healthcare provider if you have changes in your eyesight (vision) during treatment with Trulicity.

Low blood sugar (hypoglycemia). Signs and symptoms of low blood sugar may include dizziness or light-headedness, confusion or drowsiness, headache, blurred vision, slurred speech, fast heartbeat, sweating, hunger, shakiness, feeling jittery, weakness, anxiety, irritability or mood changes.

Serious allergic reactions. Stop using Trulicity and get medical help right away if you have any symptoms of a serious allergic reaction which may include: swelling of your face, lips, tongue or throat, problems breathing or swallowing, severe rash or itching, fainting or feeling dizzy, or very rapid heartbeat.

Acute kidney injury. In people who have kidney problems, diarrhea, nausea, and vomiting may cause a loss of fluids (dehydration). This may cause kidney problems to get worse.

Severe stomach problems. Trulicity may cause stomach problems, which could be severe.

Common side effects

The most common side effects of Trulicity include nausea, diarrhea, vomiting, abdominal pain and decreased appetite. These are not all the possible side effects of Trulicity.

Tell your doctor if you have any side effects. You can report side effects at 1-800-FDA-1088 or www.fda.gov/medwatch.

Before using

- Your healthcare provider should show you how to use Trulicity before you use it for the first time.
- Before you use Trulicity, talk to your doctor about low blood sugar and how to manage it.

Review these questions with your doctor:

- □ Do you have other medical conditions, including problems with your pancreas, kidneys, liver, or stomach, or have a history of diabetic retinopathy (vision problems related to diabetes)?
- Do you take other diabetes medicines, such as insulin or
- □ Do you take any other prescription medicines or over-thecounter drugs, vitamins or herbs?

Review the list below with your doctor. Trulicity may not be right for you if:

- You are pregnant or plan to become pregnant or breastfeed or plan to breastfeed.
- ☐ You have type 1 diabetes or diabetic ketoacidosis.☐ You have or have had an inflamed pancreas (pancreatitis).
- ☐ You have severe intestinal or stomach problems, such as slowed emptying or problems with digesting food.
- ☐ You are a child under 18 years old.

How to take

- Read the Instructions for Use that come with Trulicity.
- Use Trulicity exactly as your doctor says.
 Do not share your Trulicity pen, syringe or needles with
- another person.

 Do not give Trulicity to other people
- Do not give Trulicity to other people
 If you take too much Trulicity, call you
- If you take too much Trulicity, call your healthcare provider or seek medical advice promptly.

Learn more

For more information, call 1-844-TRU-INFO (1-844-878-4636) or go to www.TRULICITY.com.

This summary provides basic information about Trulicity but does not include all information known about this medicine. Read the information that comes with your prescription each time your prescription is filled. This information does not take the place of talking with your doctor. Be sure to talk to your doctor or other healthcare provider about Trulicity and how to take it. Your doctor is the best person to help you decide if Trulicity is right for you.

Trulicity® is a registered trademark owned or licensed by Eli Lilly and Company, its subsidiaries, or affiliates.

DG CONHL BS FEB 2020

PP-DG-US-3210 07/2020 ©Lilly USA, LLC 2020. All rights reserved.



^{*}Along with diet and exercise.